

No-go theorems for ψ -epistemic models based on a continuity assumption

M. K. Patra¹, S. Pironio¹, and S. Massar¹

¹*Laboratoire d'Information Quantique, Université libre de Bruxelles (ULB), 1050 Bruxelles, Belgium*

The quantum state ψ is a mathematical object used to determine the probabilities of different outcomes when measuring a physical system. Its fundamental nature has been the subject of discussions since the inception of quantum theory: is it *ontic*, that is, does it correspond to a real property of the physical system? Or is it *epistemic*, that is, does it merely represent our knowledge about the system? Assuming a natural continuity assumption and a weak separability assumption, we show here that epistemic interpretations of the quantum state are in contradiction with quantum theory. Our argument is different from the recent proof of Pusey, Barrett, and Rudolph and it already yields a non-trivial constraint on ψ -epistemic models using a single copy of the system in question.

INTRODUCTION

Quantum theory textbooks usually start from the hypothesis that to every physical system corresponds a mathematical object – a ray in Hilbert space – called the *quantum state*. They then go on to deduce the multitude of quantitative predictions that make quantum theory so successful. But does the quantum state correspond to a real physical state or does it merely represent an observer's knowledge about the underlying reality? A major reason for doubting the reality of the quantum state is that it cannot be observed directly: it can only be reconstructed indirectly by lengthy state estimation procedures [1, 2]. Furthermore, an epistemic interpretation of the quantum state could provide an intuitive explanation for many counterintuitive quantum phenomena and paradoxes, such as the measurement postulate and wavefunction collapse [3–5].

To formulate with precision the above question, we assume, following [6], that every quantum system possesses a real physical state (also called *ontic state*), denoted λ , which is independent of the observer. When a measurement is performed on the system, the probabilities to get different outcomes are determined by λ . If an ensemble of such systems is prepared, different members of the ensemble may be found in different states λ . A preparation procedure Q therefore corresponds in general to a probability distribution $P(\lambda|Q)$ over the real states. The probability to obtain the outcome r when preparation Q is followed by measurement M is $P(r|M, Q) = \sum_{\lambda} P(\lambda|Q)P(r|M, \lambda)$. Such a model will reproduce the quantum predictions if $P(r|M, Q) = \langle \psi_Q | \mathcal{M}_r | \psi_Q \rangle$, where ψ_Q is the quantum state assigned by quantum theory to the preparation Q and $\mathcal{M}_r$ is the quantum operator describing the measurement.

We can now distinguish two classes of models of the above type. A model is said to be *ψ -ontic* if the preparation of distinct pure quantum states always give rise to distinct real states. That is, for every λ either $P(\lambda|Q) = 0$ or $P(\lambda|Q') = 0$ if the preparations Q and Q' correspond to different quantum states $|\psi_Q\rangle \neq |\psi_{Q'}\rangle$. In this case, every real state λ is compatible with a unique pure quantum state. The quantum state is “encoded” in λ and we can consider it to represent a real property of the sys-

tem, akin, e.g., to the total energy of a system in classical physics [7]. In the second class of models, known as *ψ -epistemic* models, preparation of distinct pure quantum states may result in the same real state λ . Formally, there exists preparations Q and Q' corresponding to distinct quantum states $|\psi_Q\rangle \neq |\psi_{Q'}\rangle$ such that both $P(\lambda|Q) > 0$ and $P(\lambda|Q') > 0$ for some λ . In this case, the quantum state is not uniquely determined by the underlying real state and has a status analogous, e.g., to the Liouville distribution in statistical physics.

While non-trivial ψ -epistemic models exist in any fixed dimension d [8], such models are necessarily highly contrived. Indeed, Pusey, Barrett, and Rudolph (PBR) have recently shown that the predictions of ψ -epistemic models are in contradiction with quantum theory under the assumption, termed *preparation independence*, that independently prepared pure quantum states correspond to product distributions over ontic states [9]. In this work, we derive two alternative no-go theorems for ψ -epistemic models based on a natural assumption of continuity. Our approach shows that already at the level of a single system there exist strong constraints on ψ -epistemic models. Furthermore, our first no-go theorem readily translates in a simple experimental test, an implementation of which has been reported in [10] using high-dimensional attenuated coherent states of light travelling in an optical fibre.

NO-GO THEOREMS FOR ψ -EPISTEMIC MODELS

The key motivation behind our result is that ψ -epistemic models should satisfy a form of continuity. Indeed, we assign an *ontic* status to ψ if a variation of ψ necessarily implies a variation of the underlying reality λ , and we assign it an *epistemic* status if a variation of ψ does not necessarily imply a variation of λ . It is then natural to assume a form of continuity for ψ -epistemic models: a slight change of ψ induces a slight change in the corresponding ensemble of λ 's in such a way that at least some λ 's from the initial ensemble will also belong to the perturbed ensemble. We use a slightly stronger form of continuity which asserts that there are real states λ in the initial ensemble that will remain part of the per-




FIG. 1. Illustration of ψ -ontic and δ -continuous ψ -epistemic models. Depicted is the space Λ of ontic states, as well as the support of the probability distribution $P(\lambda|Q_k)$ for preparation Q_k associated to distinct pure states ψ_k , $k = 1, \dots, 5$. In ψ -ontic models (left) distinct quantum states give rise to probability distribution $P(\lambda|Q_k)$ with no overlap. In δ -continuous ψ -epistemic models (right), states that are close to each other (such as $\{\psi_1, \psi_2, \psi_3\}$ and $\{\psi_3, \psi_4, \psi_5\}$) all share common ontic states. However states that are further from each other (such as ψ_1 and $\{\psi_4, \psi_5\}$) do not necessarily have common ontic states λ .

turbed ensemble, no matter how we perturb the initial state, provided this perturbation is small enough. Models that violate this condition are presumably very contrived. Formally this continuity condition is defined as follows (see Fig. 1 for a depiction of the difference between ψ -ontic and δ -continuous ψ -epistemic models).

Definition (δ -continuity). Let $\delta > 0$ and let B_ψ^δ be the ball of radius δ centred on $|\psi\rangle$, i.e., B_ψ^δ is the set of states $|\phi\rangle$ such that $|\langle\phi|\psi\rangle| \geq 1 - \delta$. We say that a model is δ -continuous if for any preparation Q , there exists an ontic state λ (which can depend on Q) such that for all preparations Q' corresponding to quantum states $|\phi_{Q'}\rangle$ in the ball $B_{\psi_Q}^\delta$ centred on the state $|\psi_Q\rangle$, we have $P(\lambda|Q') > 0$.

Our first result is a constraint on δ -continuous models for single systems.

Theorem 1. *There are no δ -continuous models with $\delta \geq 1 - \sqrt{(d-1)/d}$ reproducing the measurement statistics of quantum states in a Hilbert space of dimension d .*

Proof. Consider d preparations Q_k ($k = 1, \dots, d$) corresponding to distinct quantum states $|\psi_k\rangle$ all contained in a ball of radius δ . By definition of a δ -continuous model, there is at least one λ for which $\min_k P(\lambda|Q_k) > 0$ and thus

$$\epsilon \equiv \sum_{\lambda} \min_k P(\lambda|Q_k) > 0. \quad (1)$$

This last quantity can be viewed as a measure of the extent to which distributions over real states overlap in the neighborhood of a given quantum state. It was also

introduced in [9] where it was shown to be related to the variational distance between the distributions $P(\lambda|Q_k)$.

Suppose now that a measurement M yielding one of the possible outcomes $r = 1, \dots, d$ is made on each of the prepared systems. A δ -continuous model then makes the prediction

$$\begin{aligned} \sum_k P(k|M, Q_k) &= \sum_k \sum_{\lambda} P(k|M, \lambda) P(\lambda|Q_k) \\ &\geq \sum_k \sum_{\lambda} P(k|M, \lambda) \min_k P(\lambda|Q_k) \\ &= \sum_{\lambda} \min_k P(\lambda|Q_k) = \epsilon > 0. \end{aligned} \quad (2)$$

According to quantum theory, however, there exist states in a Hilbert space of dimension d contained in a ball of radius $\delta = 1 - \sqrt{(d-1)/d}$ such that the left-hand side of Eq. (2) is equal to 0. To show this, let $\{|j\rangle : j = 1, \dots, d\}$ be a basis of the Hilbert space. Consider the d distinct states $|\psi_k\rangle = \frac{1}{\sqrt{d-1}} \sum_{j \neq k} |j\rangle$. These states are all at mutual distance $|\langle\psi_k|\psi\rangle| = \sqrt{(d-1)/d}$ from the state $|\psi\rangle = \frac{1}{\sqrt{d}} \sum_j |j\rangle$. Let the measurement M be the measurement in the basis $\{|j\rangle\}$. Then $P(k|M, Q_k) = 0$ for all $k = 1, \dots, d$ and thus $\sum_k P(k|M, Q_k) = 0$. (These states and measurements were considered in the $d = 3$ case in [11]). $\square$

Note that the above result also applies if we only require δ -continuity to hold around some fixed quantum states rather than for all states in Hilbert space. Interestingly, the ψ -epistemic model of [8] for Hilbert spaces of dimension d is δ -continuous around a specific state with a value of δ saturating the above bound.

Though one expects a ψ -epistemic model to be δ -continuous for some value of δ , the bound derived in Theorem 1 may a priori seem arbitrary. This motivates the following definition.

Definition (Continuity). A ψ -epistemic model is continuous if there exists a non-zero $\delta > 0$ such that it is δ -continuous.

Note that Theorem 1 implies that continuous ψ -epistemic cannot reproduce the measurement statistics of quantum states in infinite dimensional Hilbert spaces.

Our second result shows that in spaces of finite dimension d there are no ψ -epistemic models that are both continuous and which satisfy the following separability assumption. A similar condition was independently introduced in [12], where it is called “compactness”. Though weaker than the preparation independence assumption explicitly used by PBR, it is already sufficient to derive their main result.

Definition (Separability). Let Q be the preparation of a physical system yielding with non-zero probability $P(\lambda|Q) > 0$ the real state λ . A model is separable if n independent copies $Q^n = (Q, \dots, Q)$ of the preparation devices yield with non-zero probability $P(\vec{\lambda} = \lambda^n|Q^n) > 0$

a system in the joint real state $\lambda^n = (\lambda, \dots, \lambda)$, for any positive integer n .

Theorem 2. *Separable continuous ψ -epistemic models cannot reproduce the measurement statistics of quantum states in a Hilbert space of dimension $d \geq 3$.*

Proof. Consider d preparations Q_k corresponding to the d distinct states $|\phi_k\rangle = \alpha|k\rangle + \beta/\sqrt{d}\sum_{i=1}^d|i\rangle$ with $k = 1, \dots, d$, $\alpha = -\sqrt{1 - [(d-2)/(d-1)]^{1/n}}$ and $\beta = -\alpha/\sqrt{d} + \sqrt{\alpha^2/d + [(d-2)/(d-1)]^{1/n}}$. It is easily checked that these states are normalised, have mutual scalar product $|\langle\phi_k|\phi_l\rangle| = [(d-2)/(d-1)]^{1/n}$ for $k \neq l$, and are all at distance $|\langle\phi_k|\phi\rangle| = 1 - \delta_{nd}$ from the state $|\phi\rangle = \frac{1}{\sqrt{d}}\sum_{i=1}^d|i\rangle$, where $\delta_{nd} = 1 - \sqrt{[1 + (d-1)^{(n-1)/n}(d-2)^{1/n}]/d}$. In a δ_{nd} -continuous model, these states share at least a common real state λ . The separability assumption then implies that the states $|\phi_k\rangle^{\otimes n}$ also share a common real state, and thus that $\epsilon_n \equiv \sum_{\lambda} \min_k P(\lambda|Q_k^n) > 0$. By the same argument as in Theorem 1, it then follows that if a measurement M yielding one of the possible outcomes $r = 1, \dots, d$ is performed on each of these systems, the quantity $\sum_k P(k|M, Q_k^n) \geq \epsilon_n > 0$.

Note now that the d n -fold copies $|\phi_k\rangle^{\otimes n}$ have mutual scalar product $|\langle\phi_k|\phi_l\rangle^{\otimes n}| = (d-2)/(d-1)$ for $k \neq l$. There therefore exists a unitary transformation U in the subspace $S_d \subset \mathbb{C}_d^{\otimes n}$ spanned by the d states $|\phi_k^{\otimes n}\rangle$ that carries out the transformation: $U|\phi_k^{\otimes n}\rangle = |\psi_k\rangle$ where $|\psi_k\rangle = \frac{1}{\sqrt{d-1}}(\sum_{j=1}^d|j\rangle - |k\rangle)$ for some basis $\{|j\rangle\}$. The states $|\psi_k\rangle$ are identical to the states used in the proof of Theorem 1. It follows that there exists a d -outcome measurement M in $\mathbb{C}_d^{\otimes n}$ which applied on the states $|\phi_k^{\otimes n}\rangle$ gives the same statistics as the measurement in the basis $\{|j\rangle\}$ applied on the states $|\psi_k\rangle$ of Theorem 1. We can therefore find a measurement such that $P(k|M, Q_k^n) = 0$ and thus $\sum_k P(k|M, Q_k^n) = 0$ in contradiction with the prediction of a δ_{nd} -continuous separable ψ -epistemic model.

We have thus shown that one can exclude δ -continuous separable models with $\delta \geq \delta_{nd}$ for any positive integers d and n . For large n this bound behaves as $\delta \gtrsim \gamma/n$, with $\gamma = (d-1)[\log(d-2) - \log(d-1)]/(2d)$, thereby implying by taking n arbitrarily large that no ψ -epistemic model for Hilbert spaces of dimension $d \geq 3$ can satisfy both the assumptions of continuity and separability. $\square$

The contradictions between quantum theory and ψ -epistemic models presented above rely on the fact that certain quantum probabilities are exactly equal to zero. Like PBR's theorem, our arguments are robust against small deviations from these predictions, as expected for instance in an experimental implementation where noise is inevitably present. Indeed, Eq. (2) implies that the measured value $\epsilon_{\text{exp}} = \sum_k P(k|M, Q_k)$ provides an upper bound on the overlap $\epsilon = \sum_{\lambda} \min_k P(\lambda|Q_k)$ of the ontic distributions $P(\lambda|Q_k)$. A small value of ϵ_{exp} therefore translates into a strong constraint on continuous ψ -

epistemic models, since it implies that these distributions have only a small common overlap. Similarly $\epsilon_{\text{exp}}^n = \sum_k P(k|M, Q_k^n)$ in Theorem 2 upper bounds the overlap $\epsilon_n = \sum_{\vec{\lambda}} \min_k P(\vec{\lambda}|Q_k^n)$ of the n -copy joint distributions $P(\vec{\lambda}|Q_k^n)$. This last quantity can simply be related to the single-copy overlap ϵ if we further make the preparation independence assumption of PBR that joint distributions $P(\vec{\lambda}|Q_k^n) = P(\lambda_1, \dots, \lambda_n|Q_k^n) = P(\lambda_1|Q_k) \dots P(\lambda_n|Q_k)$ are product of individual distributions, which then implies $\epsilon_n = \epsilon^n$.

Note that for notational simplicity we have formulated our results in the case where the set $\Lambda = \{\lambda\}$ of real states is finite or denumerable. The generalisation of Theorems 1 and 2 to measurable spaces is given in Appendix A. This generalisation is important since reproducing the predictions of even a single qubit requires a non-countable number of real states [13–15].

DISCUSSION

In his seminal paper on the probabilistic interpretation of quantum theory, Born gave the wavefunction a functional interpretation: a mathematical object from which the probabilities of different measurement outcomes can be determined [16]. But the fundamental nature of this object, a real physical wave or a summary of our knowledge about physical systems, is a question that has divided physicists ever since. A precise formulation of these two alternatives, opening the way to clear-cut answers, was provided by Harrigan and Spekkens [6]. If the wavefunction corresponds to a real, ontic, property of physical systems, the preparation of a system in different pure quantum states should always result in different physical states. If, on the other hand, the wavefunction has an epistemic status, such preparations should sometimes result in the same underlying physical state. PBR have recently introduced a no-go theorem that, given certain assumptions, rules out this latter possibility [9], thus awarding ontic status to the wavefunction. This result can also be seen as a constraint on the structure of possible extensions or generalizations of quantum theory. If they reproduce the present quantum predictions, such theories can only *supplement* the wavefunction ψ with additional variables λ' , i.e., a system should be described by a physical state of the form $\lambda = (\psi, \lambda')$.

The theorem of PBR relies on two main assumptions. The first, which is also unquestioned in the present work, is that a system has a real and objective state λ that is independent of the observer. The second is an assumption of “preparation independence”, which states that independently prepared systems are described by independent product distributions over real states. It can be replaced by the weaker separability assumption used here.

In the present work, we reached the same conclusion as PBR using a simple argument that relies on a natural

assumption of continuity. Our notion of continuity captures the intuition that in a model where the quantum state is epistemic, a variation of ψ does not necessarily imply a variation of the underlying real state λ . We derived a fundamental limit on the degree of continuity of ψ -epistemic models, as parametrized by a quantity δ , already at the level of single quantum systems (Theorem 1). Constraints on ψ -epistemic models using single quantum systems have also been obtained independently in [17] using a different assumption termed *ontic indifference*. In Appendix B, we show how our results on δ -continuous ψ -epistemic models can recover those of [17]. Combining our continuity assumption with a separability assumption, we then showed that no ψ -epistemic model can reproduce all the predictions of quantum theory (Theorem 2).

Besides their simplicity and the fact that they already constrain ψ -epistemic models for single quantum systems, an interest of our results is that they are easy to implement experimentally. Such an experimental test based on Theorem 1 has been reported in [10]. Overall, our theoretical analysis and the experimental results presented in [10] provide further evidence for the reality of the quantum state.

Acknowledgments. M. P. would like to thank Rob Spekkens for helpful discussions. We thank anonymous referees for useful comments. We acknowledge financial support from the European Union under project QCS, from the FRS-FNRS under project DIQIP, and from the Brussels-Capital Region through a BB2B grant.

Appendix A: General formulation of the no-go theorems in the context of measurable spaces

Here we formulate our main result in the context of measurable spaces. Two main cases are of interest: when the space of ontic states Λ is discrete, and when it is absolutely continuous. The discrete case was treated in the main text. It has the advantage of notational simplicity, but is probably difficult to justify as it is known that reproducing the predictions of quantum theory requires that the space of ontic states Λ be continuous [13–15]. The absolutely continuous case is more natural, but the notation and proofs are slightly more cumbersome. In the following we first introduce a general measure theoretic framework for ontic states, and then restrict to the absolutely continuous case.

The state space Λ is assumed to be a (probability) measure space with a fixed σ -algebra Ω of events. We assume the existence of a reference measure m on Λ . For a measurable set $A \subset \Lambda$, $m(A) = \int \chi_A dm(\lambda)$, where χ_A is the indicator function for A (1 on A and 0 elsewhere).

Each preparation Q induces a conditional probability measure $P(-|Q)$ defined on Ω . From the Lebesgue decomposition theorem [18] we infer that the measure $P(-|Q)$ can be decomposed as

$$P(-|Q) = P_s(-|Q) + P_a(-|Q),$$

where the measures $P_s(-|Q)$ and $P_a(-|Q)$ are respectively singular and absolutely continuous with respect to m . Recall that a measure μ is concentrated on a set C if for any measurable set B , $\mu(B) = \mu(B \cap C)$, and that two measures are singular with respect to each other if they are concentrated on disjoint sets. Furthermore the singular part $P_s(-|Q)$ can be decomposed into a discrete singular part which consists of a sum of Dirac measures (“ δ functions”) and a singular continuous part.

In the remainder we consider the case where $P(-|Q)$ only consists of an absolutely continuous part. This is the assumption that is generally made in the literature. The case where there is only a discrete singular part was discussed in the main text. Note that our results can be extended to the case where either the absolutely continuous or the discrete singular part are non-zero. Our results do not apply to the case where there is only a singular continuous part.

We denote by S_m the set on which m is concentrated. The fact that the measure $P(-|Q)$ is absolutely continuous with respect to m ($P(A|Q) = 0$ whenever $m(A) = 0$) implies that there is an m -measurable function $p(\lambda|Q)$ on Λ such that for all $A \in \Omega$

$$P_a(A|Q) = \int_A p(\lambda|Q) dm(\lambda).$$

In integration theory the function $p(\lambda|Q)$ is called the Radon-Nikodym derivative [18].

We further suppose that the probability of obtaining result r given measurement M and ontic state λ is given by an m -measurable function $p(r|M, \lambda)$. Then we have

$$P(r|M, Q) = \int_{S_m} p(r|M, \lambda) p(\lambda|Q) dm(\lambda). \quad (\text{A1})$$

We can define the support Sup_Q of the function $p(\lambda|Q)$ as the complement of the set on which it vanishes. Since $m(Sup_Q) = m(S_m \cap Sup_Q)$ there will be no loss of generality if we restrict the support to S_m . (Note that if the measure is not absolutely continuous, we cannot introduce in general the concept of “support” of the measures $P(-|Q)$ as it requires a topology on Λ which we do not assume).

Given the above framework, we now state and prove the results presented in the main text in the case of absolutely continuous measures. This includes the case where $\Lambda = \mathbb{R}^n$ with m the standard measure on $\mathbb{R}^n$.

Definition (ψ -epistemic model). *A statistical model is ψ -epistemic if there exist two preparation procedures Q_1 and Q_2 giving two distinct quantum states such that $m(Sup_{Q_1} \cap Sup_{Q_2}) > 0$.*

Definition (Continuity). *A ψ -epistemic model is δ -continuous if, given the preparation Q of the pure quantum state ψ_Q , there is a set $A^\delta \in \Omega$ (which can depend on Q) such that for all states ϕ satisfying $1 - |\langle \phi | \psi \rangle| \leq \delta$, $A^\delta \subset Sup_{Q_\phi}$ and $\int_{A^\delta} p(\lambda|Q_\phi) dm(\lambda) > 0$. Note that this implies that $m(A^\delta) > 0$. A model is continuous, if it is δ -continuous for some $\delta > 0$.*

Why is δ -continuity a “natural” condition? Informally the reason is that –except in rare occasions– physical quantities depend continuously on their parameters. Discontinuities are generally attributable to mathematical idealisation. To illustrate this in the present case, suppose that we can write $Q = (Q', \psi_Q)$, where ψ_Q is the pure state produced by Q , and Q' are additional variables specifying the preparation. Furthermore suppose that λ, Q', ψ_Q all belong to compact subsets of real euclidean space $\mathbb{R}^n$, and that the probability distribution $p(\lambda|Q', \psi_Q)$ is a continuous function of all its parameters. Then it is uniformly continuous, and hence δ -continuous for some $\delta > 0$.

Proof of Theorem 1. Let the dimension of the quantum state space be d . We consider the same d independent non-orthogonal states $\psi_k = \frac{1}{\sqrt{d-1}} \sum_{j \neq k} |j\rangle$ as in the main text. Suppose the model is δ -continuous for $\delta \geq 1 - \sqrt{(d-1)/d}$. Then there exists a set $A^\delta \subset \text{Sup}_{\psi_k}$ for all $k = 1, \dots, d$ with $m(A^\delta) > 0$. Define a new function $\tilde{p}(\lambda) = \min_k p(\lambda|\psi_k)$. Then we have

$$\begin{aligned} \sum_k p(k|\psi_k) &= \int_{\Lambda} \sum_k p(k|\lambda) p(\lambda|\psi_k) dm(\lambda) \\ &\geq \int_{A^\delta} \sum_k p(k|\lambda) p(\lambda|\psi_k) dm(\lambda) \\ &\geq \int_{A^\delta} \sum_k p(k|\lambda) \tilde{p}(\lambda) dm(\lambda) \\ &= \int_{A^\delta} \tilde{p}(\lambda) dm(\lambda). \end{aligned} \quad (\text{A2})$$

It remains to show that $\int_{A^\delta} \tilde{p}(\lambda) dm(\lambda)$ is strictly positive. To this end we note that for each function $p(\lambda|\psi_k)$ there must be a positive integer N_k such that the set $T_{N_k} = \{\lambda \in A^\delta | p(\lambda|\psi_k) > 1/N_k\}$ satisfies $m(T_{N_k}) > 0$. Otherwise $\int_{A^\delta} p(\lambda|\psi_k) dm(\lambda) = 0$, contrary to our hypothesis. Let $M = \max\{N_k\}$. Then for the set $T_M \equiv \{\lambda \in A^\delta | \tilde{p}(\lambda) > 1/M\}$ we have $T_M \supset T_{N_k}$ for all k and hence $m(T_M) > 0$. We conclude that

$$\int_{A^\delta} \tilde{p}(\lambda) dm(\lambda) \geq \frac{1}{M} m(T_M) > 0.$$

We thus arrive at the contradiction $0 = \sum_k p(k|\psi_k) > 0$. $\square$

Definition (Separability). Consider a preparation Q of a physical system and any subset of ontic states $A \in \Lambda$ that occurs with non-zero probability $P(A|Q) > 0$. Consider n identical copies of the preparation devices, which when used yield preparation $Q^n = (Q, \dots, Q)$. Then the product set $A^n = A \times \dots \times A$ occurs with non-zero probability: $P(A^n|Q^n) > 0$.

Note that according to this definition the space of ontic states for n independent preparations may be larger than the product $\Lambda \times \dots \times \Lambda$. This does not affect the definition of separability.

Proof of Theorem 2. The proof of Theorem 2 in the case of absolutely continuous measures is a simple combination of the arguments given in the main text in the case of discrete singular measures, and the proof of Theorem 1 for the case of absolutely continuous measures. $\square$

Appendix B: Relation between our continuity assumption and ontic indifference

Hardy recently proposed a no-go theorem for ψ -epistemic models for a single system that satisfies *ontic indifference*, or *restricted ontic indifference* (see definitions below) [17]. Here we show that ontic indifference is an extremely strong condition for ψ -epistemic models, since it implies that there are ontic states λ that are shared by all quantum states in the Hilbert space. It is therefore immediately in contradiction with the predictions of quantum theory since this implies that orthogonal quantum states have overlapping ontic distributions. Restricted ontic indifference does not suffer from this problem, but on the other hand it breaks the invariance of Hilbert space by promoting one state to a specific status. We show how the proof methods we use to rule out δ -continuous ψ -epistemic models can be easily transposed to provide no-go results for ψ -epistemic models obeying restricted ontic indifference.

Definition (Ontic indifference [17]). Any quantum transformation on a system which leaves unchanged any given pure state, $|\psi\rangle$, can be performed in such a way that it does not affect the underlying ontic states $\lambda \in \Lambda_{|\psi\rangle}$ in the ontic support of that pure state. By the ontic support of a given state, $|\psi\rangle$, one means the set $\Lambda_{|\psi\rangle}$ of ontic states λ which might be prepared when the given pure state is prepared.

Observation. Ontic indifference implies the following fact: if two distinct states $|\psi\rangle \neq |\psi'\rangle$ share the same ontic state λ , that is $\lambda \in \Lambda_{|\psi\rangle}$ and $\lambda \in \Lambda_{|\psi'\rangle}$, and if U is a unitary transformation that leaves $|\psi\rangle$ invariant, $U|\psi\rangle = |\psi\rangle$, then $\lambda \in \Lambda_{U|\psi'\rangle}$ belongs to the ontic support of $U|\psi'\rangle$.

This follows from considering the following 4 preparations: a specific preparation P_1 of $|\psi\rangle$, the preparation P_1 followed by unitary U (that leaves $|\psi\rangle$ invariant), a specific preparation P_2 of $|\psi'\rangle$, and the preparation P_2 followed by unitary U (which prepares $U|\psi'\rangle$). Then we note that preparation P_1 sometimes yields the ontic state λ , that U leaves λ invariant, that preparation P_2 sometimes yields the ontic state λ , hence preparation P_2 followed by U sometimes yields the ontic state λ , and therefore $\lambda \in \Lambda_{U|\psi'\rangle}$.

This observation has the following consequence, which implies directly that no ψ -epistemic model can reproduce the predictions of quantum theory (since it implies in particular that orthogonal quantum states share a common real state λ).

Theorem 3. *If any two distinct states $|\psi_1\rangle \neq |\psi_2\rangle$ share the same ontic state λ , that is $\lambda \in \Lambda_{|\psi_1\rangle}$ and $\lambda \in \Lambda_{|\psi_2\rangle}$, and if ontic indifference holds, then all states in the Hilbert space share the ontic state λ , that is $\lambda \in \Lambda_{|\psi\rangle}$ for all $|\psi\rangle$.*

Proof. To prove this consider the following sets S_k ($k \geq 0$) of states, where $S_0 = \{|\psi_1\rangle, |\psi_2\rangle\}$ and $S_k = \{|\psi\rangle : \text{there exist } |\phi\rangle, |\phi'\rangle \in S_{k-1} \text{ and a unitary } U \text{ such that } U|\phi\rangle = |\phi\rangle, U|\phi'\rangle = |\psi\rangle\}$. That is, S_k is the set of states that can be obtained from S_{k-1} by a unitary that leaves one of the states in S_{k-1} invariant.

By hypothesis, both states in S_0 share the same ontic state λ . Hence by the above observation, all states in S_k , for all $k = 0, 1, 2, \dots$, also share the state λ . Now consider the case of a 2-dimensional Hilbert space. If $|\langle\psi_1|\psi_2\rangle| = \cos\theta$ are at an angle θ from each other (as measured on the Bloch sphere), then it is easy to show (just draw the successive sets S_k) that the size of S_k increases exponentially, such that when $k \geq O(\log\theta)$, S_k comprises the whole Bloch sphere. The case of a 2-dimensional Hilbert space helps to visualize what is happening, but is not a restriction, since the linear span of the set S_1 is the whole Hilbert space. $\square$

We now show how a simple application of our results provides a no-go theorem for ψ -epistemic models satisfying restricted ontic indifference.

Definition (Restricted ontic indifference [17]). *Any quantum transformation on a system which leaves un-*

changed the pure state $|\psi\rangle$ can be performed in such a way that it does not affect the underlying ontic states, $\lambda \in \Lambda_{|\psi\rangle}$, in the ontic support of that pure state.

Theorem 4. *The quantum predictions are incompatible with ψ -epistemic models satisfying the assumptions of restricted ontic indifference and separability, and such that there exists a state $|\psi'\rangle \neq |\psi\rangle$ sharing a common ontic state λ with $|\psi\rangle$, where $|\psi\rangle$ is the particular state in the definition of restricted ontic indifference.*

Proof. Let $d \geq 2$ be an integer such that $|\langle\psi'|\psi\rangle| \leq \sqrt{\frac{d-1}{d}}$. Add an ancilla $|0\rangle$ to the space such that the dimension of the full Hilbert space is greater or equal to d . By the assumption of separability the states $|\Psi\rangle = |\psi\rangle|0\rangle$ and $|\Psi'\rangle = |\psi'\rangle|0\rangle$ share a common ontic state λ . Choose a basis such that $|\Psi\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |i\rangle$, and coefficients $\alpha_i \in \mathbb{R}$, $i = 1, \dots, d$ such that $\alpha_1 = 0$, $\sum_{i=2}^d \alpha_i^2 = 1$, $\frac{1}{\sqrt{d}} \sum_{i=2}^d \alpha_i = |\langle\Psi'|\Psi\rangle|$ (this is always possible since $|\langle\Psi'|\Psi\rangle| \leq \sqrt{\frac{d-1}{d}}$). Define the states $|\Phi_k\rangle = \sum_{i=1}^d \alpha_{i-k} |i\rangle$. (Note that these states coincide with the states Ψ_k used in the main text if $\alpha_i = \frac{1}{\sqrt{d-1}}$, $i = 2, \dots, d$). The Observation above implies that all the states $|\Phi_k\rangle$ share the ontic state λ , since $|\langle\Phi_k|\Psi\rangle| = |\langle\Psi'|\Psi\rangle|$. We then obtain a contradiction with the predictions of quantum theory exactly as in the proof of Theorem 1 given in the main text. $\square$

-
- [1] M. G. A. Paris and J. Rehacek (Eds.), *Quantum State Estimation*, Lect. Notes Phys. **649**, 1, Springer (2004).
 - [2] Y. Aharonov and L. Vaidman, *Measurement of the Schrödinger wave of a single particle*, Phys. Lett. A **178**, 38 (1993).
 - [3] C. M. Caves, C. A. Fuchs, and R. Schack, *Quantum probabilities as Bayesian probabilities*, Phys. Rev. A **65**, 022305 (2002).
 - [4] C. Brukner and A. Zeilinger, *Information and fundamental elements of the structure of quantum theory*, in “Time, Quantum, Information”, edited by L. Castell and O. Ischebeck, Springer (2003).
 - [5] R. W. Spekkens, *Evidence for the epistemic view of quantum states: A toy theory*, Phys. Rev. A **75**, 032110 (2007).
 - [6] N. Harrigan and R. W. Spekkens, *Einstein, Incompleteness, and the Epistemic View of Quantum States*, Found. Phys. **40**, 125 (2010).
 - [7] M. Leifer, *PBR, EPR, and all that jazz*, The Quantum Times **6**(3) (2011).
 - [8] P. G. Lewis, D. Jennings, J. Barrett, and T. Rudolph, *Distinct Quantum States Can Be Compatible with a Single State of Reality*, Phys. Rev. Lett. **109**, 150404 (2012).
 - [9] M. F. Pusey, J. Barrett, and T. Rudolph, *On the reality of the quantum state*, Nat. Phys. **8**, 476 (2012).
 - [10] M. K. Patra, L. Olislager, F. Duport, J. Safioui, S. Pironio, S. Massar, arXiv:1211.1179v1
 - [11] C. M. Caves, C. A. Fuchs, and R. Schack, *Conditions for compatibility of quantum-state assignments*, Phys. Rev. A **66**, 062111 (2002).
 - [12] M. Schlosshauer and A. Fine, *Implications of the Pusey-Barrett-Rudolph Quantum No-Go Theorem*, Phys. Rev. Lett. **108**, 260404 (2012).
 - [13] L. Hardy, *Quantum ontological excess baggage*, Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics **35**, 267 (2004).
 - [14] A. Montina, *Exponential complexity and ontological theories of quantum mechanics*, Phys. Rev. A **77**, 022104 (2008).
 - [15] A. Montina, *State-space dimensionality in short-memory hidden-variable theories*, Phys. Rev. A **83**, 032107 (2011).
 - [16] M. Born, *Zur Quantenmechanik der Stoßvorgänge*, Zeitschrift für Physik **37**, 863 (1926).
 - [17] L. Hardy, *Are quantum states real?*, arXiv:1205.1439 (2012).
 - [18] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, New York (1987).